

## 蔬菜蟲害防治曆

| 作物  | 主要蟲害   | 藥劑                 | 稀釋倍數(倍) | 安全採收(天) |
|-----|--------|--------------------|---------|---------|
| 蔬菜  | 蔬菜小金花蟲 | 43%佈飛松乳劑           | 1,000   | 12      |
|     |        | 2.8%賽洛寧乳劑          | 1,000   | 12      |
|     |        | 50%免速達可濕性粉劑        | 1,000   | 15      |
| 青蔥  | 甜菜夜蛾   | 19.7%得芬諾水懸劑        | 2,000   | 9       |
|     |        | 2.8%畢芬寧乳劑          | 666     | 10      |
|     |        | 4.4%綠芬隆乳劑          | 1,500   | 10      |
|     |        | 9.6%氟芬隆水分散性乳劑      | 3,000   | 12      |
|     | 韭馬     | 50%馬拉松乳劑           | 1,500   | 4       |
|     |        | 40%納乃得水溶性粒劑        | 1,500   | 10      |
|     |        | 50%培丹水溶性粉劑         | 1,000   | 15      |
|     |        | 50%賽達松乳劑           | 1,000   | 15      |
| 沖香瓜 | 番茄斑蟊蛾  | 2%阿巴汀乳劑            | 2,000   | 6       |
|     |        | 75%賽滅淨可濕性粉劑        | 7,500   | 15      |
|     | 棉蚜     | 40.4%賽果培溶液         | 4,000   | 21      |
|     | 南黃翅馬   | 10%克凡派水懸劑          | 1,000   | 6       |
|     | 銀葉粉蝨   | 20%達特南水溶性粒劑        | 3,000   | 6       |
|     |        | 10%賽速安水溶性粒劑        | 4,000   | 6       |
|     |        | 2.8%畢芬寧乳劑          | 1,000   | 6       |
|     | 黃守瓜    | 50%加保利可濕性粉劑        | 1,000   | 7       |
|     | 瓜實蠅    | 25%馬拉松可濕性粉劑+蛋白質水解物 | 400     | 7       |
|     | 二點葉蟊   | 1%密滅汀乳劑            | 1,500   | 6       |
|     | 甜菜夜蛾   | 5%因滅汀可溶性粒劑         | 4,000   | 6       |
| 番茄  | 銀葉粉蝨   | 20%亞滅培水溶性粒劑        | 4,000   | 3       |
|     | 夜蛾類    | 14.5%因得克水懸劑        | 2,000   | 6       |
|     |        | 2.15%因滅汀乳劑         | 1,500   | 6       |
|     |        | 34%硫敵克水懸劑          | 1,000   | 3       |
|     |        | 4.4%綠芬隆乳劑          | 1,500   | 6       |
|     |        | 44.9%聯斯松乳劑         | 1,000   | 6       |
|     | 番茄斑蟊蛾  | 2%阿巴汀乳劑            | 1,000   | 12      |
|     |        | 75%賽滅淨可濕性粉劑        | 4,000   | 12      |
| 南瓜  | 瓜蟻     | 50%加保利可濕性粉劑        | 1,000   | 7       |
|     | 蚜蟲     | 25%滅滅淨可濕性粉劑        | 2,000   | 12      |
|     | 黃守瓜    | 50%加保利可濕性粉劑        | 1,000   | 7       |
|     | 瓜實蠅    | 25%馬拉松可濕性粉劑+蛋白質水解物 | 400     | 7       |
| 胡瓜  | 二點葉蟊   | 1%密滅汀              | 1,500   | 3       |
|     | 南黃翅馬   | 4.95%芬普尼水懸劑        | 2,000   | 6       |
|     |        | 9.6%益蟲腺液           | 1,500   | 6       |

| 作物         | 主要蟲害    | 藥劑           | 稀釋倍數(倍) | 安全採收(天) |
|------------|---------|--------------|---------|---------|
| 甜瓜         | 銀葉粉蝨    | 0.5%可尼丁粒劑    |         |         |
|            | 蚜蟲      | 10%無尼胺水分散粒劑  | 4,000   | 3       |
|            |         | 50%滅滅淨水分散粒劑  | 4,000   | 12      |
|            |         | 25%滅滅淨可濕性粉劑  | 2,000   | 12      |
|            | 黃守瓜     | 50%加保利可濕性粉劑  | 1,000   | 7       |
| 菜豆<br>(蠶豆) | 根腐病、莖腐病 | 5%二硫松粒劑      | 1克/2穴   |         |
|            |         | 10%福瑞松粒劑     | 1克/4穴   |         |
|            | 蚜蟲      | 40%納乃得可濕性粉劑  | 1,500   | 10      |
|            |         | 3.75%寧滅寧乳劑   | 1,000   | 12      |
|            | 小翅葉蟬    | 25%丁基加保可濕性粉劑 | 1,500   | 15      |
|            | 眼蠅目幼蟲   | 3%蘇力菌可濕性粉劑   | 700     | 9       |
|            |         | 58%乃力松乳劑     | 1,000   | 4       |
|            |         | 50%馬拉松乳劑     | 500     | 4       |
|            | 葉蟥      | 2.8%賽洛寧乳劑    | 2,000   | 10      |
|            | 薊馬      | 2.8%畢芬寧乳劑    | 1,000   | 3       |
|            |         | 2.8%賽洛寧乳劑    | 1,000   | 3       |
|            |         | 10%美文松乳劑     | 300     | 3       |
|            | 番茄斑潛蛾   | 75%賽滅淨可濕性粉劑  | 5,000   | 6       |
|            |         | 2.8%賽洛寧乳劑    | 1,000   | 6       |
|            | 豆夾螟     | 50%加保利可濕性粉劑  | 1,000   | 7       |
| 綠竹筴        | 蚜蟲      | 5%可尼丁粒劑      | 9公斤/分地  | 3       |
|            |         | 25%滅滅松乳劑     | 1:4     | 18      |
|            | 盲椿象     | 30%撲芬松乳劑     | 1,500   | 18      |
|            |         | 25%滅滅松乳劑     | 1:4     | 18      |
|            |         | 20%芬化利乳劑     | 2,000   | 未訂      |
|            | 捲葉蛾     | 5.7%賽扶寧乳劑    | 250     | 9       |
| 甜椒         | 紅蜘蛛     | 新殺蟎          | 500     | 15      |
|            | 二點葉蟥    | 25%畢達本可濕性粉劑  | 3,000   | 3       |
|            |         | 1%密滅汀乳劑      | 1,500   | 3       |
|            |         | 2.8%畢芬寧乳劑    | 1,500   | 6       |
|            |         | 2.5%賽洛寧微乳劑   | 1,000   | 6       |
|            |         | 2.8%賽洛寧乳劑乳劑  | 1,000   | 3       |
|            | 薊馬      | 9.6%益達胺溶液    | 2,000   | 6       |
|            |         | 2.8%第滅寧水基乳劑  | 1,000   | 12      |
|            |         | 2.8%第滅寧乳劑    | 1,000   | 12      |



| 作物  | 主要蟲害   | 藥劑                  | 稀釋倍數(倍) | 安全採收(天) |
|-----|--------|---------------------|---------|---------|
| 甜椒  | 二點小綠葉蟬 | 2.8%畢芬寧乳劑           | 1,000   | 6       |
|     | 蚜蟲     | 2.8%畢芬寧乳劑           | 1,500   | 15      |
|     | 夜蛾類    | 4.4%綠芬隆乳劑           | 1,500   | 6       |
|     |        | 2.8%畢芬寧乳劑           | 1,500   | 10      |
| 西瓜  | 番茄斑蟊蛾  | 2%阿巴汀乳劑             | 2,000   | 6       |
|     |        | 75%賽滅淨可濕性粉劑         | 7,500   | 15      |
|     | 棉蚜     | 25%丁基加保扶可濕性粉劑       | 700     | 15      |
|     | 南黃薊馬   | 20%亞滅培水溶性粉劑         | 4,000   | 6       |
|     |        | 9.6%益達胺溶液           | 1,500   | 6       |
|     |        | 2.8%賽格寧乳劑           | 2,000   | 7       |
|     | 黃守瓜    | 50%加保利可濕性粉劑         | 1,000   | 7       |
|     | 瓜實蠅    | 25%馬拉松可濕性粉劑+蛋白質水解物  | 400     | 7       |
|     | 二點葉蟊   | 1.5%亞蘇蟊水懸劑          | 1,500   | 6       |
|     |        | 10%依殺蟊水懸劑           | 4,000   | 6       |
|     |        | 1%密滅汀乳劑             | 1,500   | 6       |
|     | 二點葉蟊   | 1.0%密滅汀乳劑           | 1,500   | 3       |
| 苦瓜  | 薊馬     | 4.95%芬普尼水懸劑         | 2,000   | 6       |
|     |        | 9.6%益達胺溶液           | 1,500   | 6       |
|     | 蚜蟲     | 25%賽滅淨可濕性粉劑         | 2,000   | 12      |
|     | 黃守瓜    | 50%加保利可濕性粉劑         | 1,000   | 7       |
|     | 瓜實蠅    | 25%馬拉松可濕性粉劑+蛋白質水解物  | 400     | 7       |
|     | 夜蛾類    | 蘇力菌水分散粒劑 32000IU/mg | 1,000   |         |
| 萬壽瓜 | 夜蛾類    | 19.7%得芬諾水懸劑         | 2,000   | 12      |
|     |        | 10%克凡派水懸劑           | 1,000   | 12      |
|     |        |                     |         |         |
|     | 薊馬     | 4.95%芬普尼水懸劑         | 2,000   | 6       |
|     |        | 9.6%益達胺水溶液          | 1,500   | 6       |
|     | 瓜實蠅    | 50%加保利可濕性粉劑         | 1,000   | 7       |
|     | 蚜蟲     | 25%賽滅淨可濕性粉劑         | 2,000   | 12      |
|     | 黃守瓜    | 50%加保利可濕性粉劑         | 1,000   | 7       |
| 白豆  | 黃條葉蟊   | 25%馬拉松可濕性粉劑+蛋白質水解物  | 400     | 7       |
|     |        | 4.95%芬普尼水懸劑         | 2,000   | 6       |
|     |        | 10%美文松乳劑            | 850     | 3       |
|     |        | 20%達特南水溶性粒劑         | 3,000   | 3       |
|     | 黃條葉蟊   | 50%培丹可溶性粉劑          | 1,000   | 10      |
| 白豆  | 黃條葉蟊   | 43%佈飛松乳劑            | 1,000   | 10      |
|     |        |                     |         |         |
|     |        |                     |         |         |
|     |        |                     |         |         |

| 作物         | 主要蟲害  | 藥 劑            | 稀釋倍數(倍)     | 安全採收(天) |
|------------|-------|----------------|-------------|---------|
| 白菜         | 小飛蛾   | 10%蘇力菌水分散性粒劑   | 2,000       |         |
|            |       | 21.5%因滅汀乳劑     | 2,000       | 6       |
|            |       | 2.8%賽洛寧乳劑      | 2,000       | 9       |
|            |       | 2.8%畢芬寧乳劑      | 2,000       | 9       |
|            |       | 10%克凡派水懸劑      | 1,000       | 10      |
|            | 蚜蟲    | 18.2%益達胺水懸劑    | 8,000       | 9       |
|            |       | 50%培丹水溶性粉劑     | 1,000       | 10      |
|            |       | 2.8%第滅寧水基乳劑    | 1,000       | 6       |
|            | 潛蛾    | 2%阿巴汀乳劑        | 2,000       | 6       |
|            |       | 8.9%賽滅淨溶液      | 800         | 7       |
|            |       | 2.8%第滅寧乳劑      | 1,000       | 6       |
|            | 斜紋夜盜蟲 | 2.8%畢芬寧乳劑      | 1,000       | 9       |
|            |       | 37.5%汰芬諾克水分性粒劑 | 1,000       | 14      |
| 甘藍<br>結球白菜 | 黃條葉蟥  | 10%美文松乳劑       | 350         | 3       |
|            |       | 20%達特南水溶性粒劑    | 3,000       | 8       |
|            |       | 50%培丹可溶性粉劑     | 1,000       | 10      |
|            |       | 43%佈飛松乳劑       | 1,000       | 10      |
|            | 菜心蛾   | 3%蘇力菌可濕性粉劑     | 1,500       |         |
|            |       | 10%美文松乳劑       | 350         | 3       |
|            | 小飛蛾   | 10%蘇力菌水分散性粒劑   | 2,000       |         |
|            |       | 2.8%賽洛寧乳劑      | 2,000       | 9       |
|            |       | 21.5%因滅汀乳劑     | 2,000       | 18      |
|            | 蚜蟲    | 10%美文松乳劑       | 350         | 8       |
|            |       | 2.8%第滅寧水基乳劑    | 1,000       | 6       |
|            |       | 18.2%益達胺水懸劑    | 8,000       | 9       |
|            | 粉虱    | 25%布芬淨可濕性粉劑    | 1,500       | 6       |
|            |       | 2.8%畢芬寧乳劑      | 1,000       | 9       |
|            |       | 9.6%益達胺溶液      | 1,000-1,500 | 9       |
|            | 潛蛾    | 2%阿巴汀乳劑        | 2,000       | 6       |
|            |       | 8.9%賽滅淨溶液      | 800         | 7       |
|            |       | 2.8%第滅寧乳劑      | 1,000       | 6       |
|            | 斜紋夜盜蟲 | 2.8%畢芬寧乳劑      | 1,000       | 9       |
|            |       | 37.5%汰芬諾克水分性粒劑 | 1,000       | 14      |
| 芥藍         | 黃條葉蟥  | 10%美文松乳劑       | 350         | 8       |
|            |       | 20%達特南水溶性粒劑    | 3,000       | 8       |
|            |       | 50%培丹可溶性粉劑     | 1,000       | 10      |
|            |       | 43%佈飛松乳劑       | 1,000       | 10      |



| 作物         | 主要蟲害    | 藥劑             | 稀釋倍數(倍)     | 安全採收(天) |
|------------|---------|----------------|-------------|---------|
| 芥藍         | 小菜蛾     | 10%蘇力菌水分散性粒劑   | 2,000       |         |
|            |         | 21.5%啞滅汀乳劑     | 2,000       | 6       |
|            |         | 2.8%賽洛寧乳劑      | 2,000       | 9       |
|            |         | 2.8%畢芬寧乳劑      | 2,000       | 9       |
|            |         | 10%克凡派水懸劑      | 1,000       | 10      |
|            | 蚜蟲      | 18.2%益達胺水懸劑    | 8,000       | 9       |
|            |         | 50%培丹水溶性粉劑     | 1,000       | 10      |
|            |         | 2.8%第滅寧水基乳劑    | 1,000       | 6       |
|            | 粉虱      | 25%布芬淨可濕性粉劑    | 1,500       | 6       |
|            |         | 2.8%畢芬寧乳劑      | 1,000       | 9       |
|            |         | 9.6%益達胺溶液      | 1,000-1,500 | 9       |
|            | 薊馬      | 2%阿巴汀乳劑        | 2,000       | 6       |
|            |         | 8.9%賽滅淨溶液      | 800         | 7       |
|            |         | 2.8%第滅寧乳劑      | 1,000       | 6       |
|            | 斜紋夜盜蟲   | 2.8%畢芬寧乳劑      | 1,000       | 9       |
|            |         | 37.5%汰芬諾克水分性粒劑 | 1,000       | 14      |
| 長豇豆        | 根蠹蛾、莖蠹蛾 | 75%賽滅淨可濕性粉劑    | 5,000       | 6       |
|            |         | 2.8%賽洛寧乳劑      | 2,000       | 10      |
|            | 豆蚜      | 58%乃力松乳劑       | 1,000       | 4       |
|            |         | 2.8%第滅寧乳劑      | 1,000       | 12      |
|            | 小綠菜蛾    | 2.8%賽洛寧乳劑      | 2,000       | 10      |
|            | 鱗翅目幼蟲   | 3%蘇力菌可濕性粉劑     | 1,500       | 1       |
|            |         | 58%乃力松乳劑       | 1,000       | 4       |
|            |         | 50%馬拉松乳劑       | 500         | 4       |
|            | 葉蟥      | 2.8%賽洛寧乳劑      | 2,000       | 10      |
|            | 薊馬      | 10%百滅寧乳劑       | 3,000       | 3       |
|            |         | 2.8%賽洛寧乳劑      | 2,000       | 10      |
|            |         | 2.8%第滅寧乳劑      | 1,000       | 12      |
|            | 番茄斑蟊    | 75%賽滅淨可濕性粉劑    | 5,000       | 6       |
|            |         | 2.8%賽洛寧乳劑      | 2,000       | 10      |
| 花椰菜<br>青花菜 | 黃條菜蟲    | 10%美文松乳劑       | 350         | 3       |
|            |         | 20%達特南水溶性粒劑    | 3,000       | 3       |
|            |         | 50%培丹可溶性粉劑     | 1,000       | 10      |
|            |         | 43%佈飛松乳劑       | 1,000       | 10      |
|            | 菜心螟     | 3%蘇力菌可濕性粉劑     | 1,500       |         |
|            |         | 10%美文松乳劑       | 850         | 3       |

| 作物         | 主要蟲害  | 藥劑             | 稀釋倍數(倍)     | 安全採收(天) |
|------------|-------|----------------|-------------|---------|
| 花椰菜<br>青花菜 | 小菜蛾   | 10%蘇力菌水分散性粒劑   | 2,000       |         |
|            |       | 2.8%賽洛寧乳劑      | 2,000       | 9       |
|            |       | 21.5%因滅汀乳劑     | 2,000       | 18      |
|            | 蚜蟲    | 10%美文松乳劑       | 850         | 3       |
|            |       | 2.8%第滅寧水基乳劑    | 1,000       | 6       |
|            |       | 18.2%益達胺水懸劑    | 8,000       | 9       |
|            | 粉虱    | 25%布芬淨可濕性粉劑    | 1,500       | 6       |
|            |       | 2.8%畢芬寧乳劑      | 1,000       | 9       |
|            |       | 9.6%益達胺溶液      | 1,000-1,500 | 9       |
|            | 薊馬    | 2%阿巴汀乳劑        | 2,000       | 6       |
|            |       | 8.9%賽滅淨溶液      | 800         | 7       |
|            |       | 2.8%第滅寧乳劑      | 1,000       | 6       |
|            | 斜紋夜盜蟲 | 2.8%畢芬寧乳劑      | 1,000       | 9       |
|            |       | 37.5%汰芬諾克水分性粒劑 | 1,000       | 14      |
| 蘿蔔         | 黃條葉蚤  | 10%歐殺滅溶液       | 200         | 6       |
|            |       | 48%佈飛松乳劑       | 1,000       | 10      |
|            |       | 2%阿巴汀乳劑        | 2,000       | 15      |
| 胡蘿蔔        | 夜蛾類   | 48.1%蘇力菌水分散性粒劑 | 1,000       |         |
|            |       | 10%克凡派水懸劑      | 1,000       | 3       |
|            |       | 2.8%賽洛寧乳劑      | 2,000       | 8       |
| 包心<br>芥菜   | 黃條葉蚤  | 10%美文松乳劑       | 350         | 3       |
|            |       | 20%達特南水溶性粒劑    | 3,000       | 3       |
|            |       | 50%培丹可溶性粉劑     | 1,000       | 10      |
|            |       | 43%佈飛松乳劑       | 1,000       | 10      |
|            | 菜心螟   | 3%蘇力菌可濕性粉劑     | 1,500       |         |
|            |       | 10%美文松乳劑       | 350         | 3       |
|            | 小菜蛾   | 10%蘇力菌水分散性粒劑   | 2,000       |         |
|            |       | 2.8%賽洛寧乳劑      | 2,000       | 9       |
|            |       | 10%克凡派水懸劑      | 1,000       | 14      |
|            |       | 21.5%因滅汀乳劑     | 2,000       | 18      |
|            | 蚜蟲    | 10%美文松乳劑       | 850         | 3       |
|            |       | 2.8%第滅寧水基乳劑    | 1,000       | 6       |
|            |       | 18.2%益達胺水懸劑    | 8,000       | 9       |
|            |       | 50%培丹水溶性粉劑     | 1,000       | 10      |
|            | 粉虱    | 25%布芬淨可濕性粉劑    | 1,500       | 6       |
|            |       | 2.8%畢芬寧乳劑      | 1,000       | 9       |
|            |       | 9.6%益達胺溶液      | 1,000-1,500 | 9       |



| 作物       | 主要蟲害  | 藥劑                 | 稀釋倍數(倍) | 安全採收(天) |
|----------|-------|--------------------|---------|---------|
| 包心<br>青菜 | 蚜蟲    | 2%阿巴汀乳劑            | 2,000   | 6       |
|          |       | 8.9%賽滅淨溶液          | 800     | 7       |
|          |       | 2.8%第滅寧乳劑          | 1,000   | 6       |
|          | 斜紋夜盜蟲 | 2.8%畢芬寧乳劑          | 1,000   | 9       |
|          |       | 37.5%汰芬諾克水分性粒劑     | 1,000   | 14      |
| 黃芽菜      | 夜蛾    | 48.1%蘇力菌水分散粒劑      | 2,000   | 9       |
|          |       | 2.8%畢芬寧乳劑          | 666     | 10      |
|          |       | 4.4%綠芬隆乳劑          | 1,500   | 10      |
|          |       | 9.6%氟芬隆水分散性乳劑      | 3,000   | 12      |
|          | 韭馬    | 50%馬拉松乳劑           | 1,500   | 4       |
|          |       | 40%納乃得水溶性粒劑        | 1,500   | 10      |
|          |       | 50%培丹水溶性粉劑         | 1,000   | 15      |
|          |       | 50%賽達松乳劑           | 1,000   | 15      |
| 大蔴       | 根蟻    | 10%敵殺滅溶液           | 1,000   | 26      |
|          | 甜菜夜蛾  | 19.7%得芬諾水懸劑        | 2,000   | 9       |
|          |       | 2.8%畢芬寧乳劑          | 666     | 10      |
|          |       | 4.4%綠芬隆乳劑          | 1,500   | 10      |
|          |       | 9.6%氟芬隆水分散性乳劑      | 3,000   | 12      |
|          | 韭馬    | 50%馬拉松乳劑           | 1,500   | 4       |
|          |       | 40%納乃得水溶性粒劑        | 1,500   | 10      |
|          |       | 50%培丹水溶性粉劑         | 1,000   | 15      |
|          |       | 50%賽達松乳劑           | 1,000   | 15      |
| 絲瓜       | 葉蟬    | 1%密滅汀乳劑            | 1,500   | 3       |
|          | 韭馬    | 4.95%芬替尼水懸劑        | 2,000   | 6       |
|          |       | 9.8%益達胺溶液          | 1,500   | 6       |
|          | 瓜蚜    | 50%加保利可濕性粉劑        | 1,000   | 7       |
|          | 蚜蟲    | 25%滅滅淨可濕性粉劑        | 2,000   | 12      |
|          | 黃守瓜   | 50%加保利可濕性粉劑        | 1,000   | 7       |
|          | 瓜實蝨   | 25%馬拉松可濕性粉劑+蛋白質水解物 | 400     | 7       |
|          | 夜蛾    | 4.95%芬替尼水懸劑        | 2,000   | 6       |



小菜蛾



切根蟲



斜紋夜盜蟲



紋白蝶



菜心爛



菜蚜



黃條葉蚤



黑守瓜



小黃瓜棉蚜



瓜蚜



花薊馬



瓜實蠅



苦瓜棉蚜



菠菜葉蟬



小白紋毒蛾



白尾紅火蚜



赤葉蟬



金線龜甲金花蟲



黑羽蛾



黃守瓜



銀葉粉蝨



小銀葉蟬



盲椿象



番茄斑潛蛾



茶細蟬



棉蚜



番茄夜蛾



茄薊馬